

Gyno-Pevaryl® 50

Local treatment of vulvo-vaginal mycoses

Properties

GYNO-PEVARYL 50 is a highly efficacious anti-mycotic for vulvo-vaginal administration. Its range of application includes the fungi occurring in the vulva and the vagina, especially *Candida albicans*.

GYNO-PEVARYL 50 makes it possible to cure cases which previously have been therapy-resistant.

GYNO-PEVARYL 50 is well tolerated locally and removes the concomitant vaginal symptoms, such as burning and itching, within a brief period of time. It does not stain the underwear.

For the treatment of partners (*Balanitis mycetica*) the product is available in cream form

Active substance

Econazole nitrate 1%

1-2-(2,4-Dichlorophenyl)-2-(4-chlorobenzyloxy)-ethylimidazole-nitrate.

Indications

Vulvo-vaginal mycoses
Mycotic balanitis

Contraindications

none of record

Dosage and application

(unless otherwise prescribed by the physician)

During a period of 2 weeks introduce one applicator charge of GYNO-PEVARYL 50 deeply into the vagina, optimally in a supine position, just before retiring for the night.

The treatment must be continued even after the disappearance of subjective complaints (itching, discharge).

Treatment of partner:

During a period of 2 weeks carefully wash glands and prepuce (foreskin) with warm water, then apply GYNO-PEVARYL 50 cream.

How to apply:

1. Open tube and screw applicator on the tube.
2. Hold tube with applicator upside in an inclined position and press cream from tube into the applicator. If the white interior part of the applicator does not move upwards automatically, loosen it with a slight pull.
3. Take a supine position, introduce applicator deeply into the vagina and empty applicator by pressing the white piston completely into the applicator.
4. Withdraw applicator.

CAUTION

Do not hold applicator by the white piston when withdrawing but by the cylinder.

Cleaning of applicator

After each application, unscrew cap of white piston and take applicator apart. Wash in luke warm water with detergent. Rinse thoroughly. Do not boil.

Supplied


GYNO-PEVARYL 50 Vaginal Cream 1%. Tube containing 40 g with applicator.

GYNO-PEVARYL 50 Vaginal Ovules, 15 ovules of 50 mg each.

Store in a refrigerator for 30 minutes prior to use.


Dispensed in pharmacies on doctor's prescriptions.

Directions for use



1. To open the tube unscrew the cap. Then pierce the seal of the tube by means of the pin on the top of the cap.

Replace the cap by the applicator.



2. Push on end of tube to bring the cream into applicator. If the piston shows resistance, pull gently. The applicator should completely be filled, unless the practising physician prescribes otherwise.

3. Remove applicator from tube. Close tube instantly with care.

4. While lying down, knees bent and spread out, insert applicator into vagina as deeply as possible. Press piston completely.

Then remove the applicator.

directions for cleaning the applicator

Pull the 2 parts apart and wash them with luke warm water and soap. Dry them and assemble the applicator by pushing the inner part back into the outer part. If the doctor specifically recommends sterilisation of the applicator, his directions for use should be followed. Never use water that is hotter than 50°C and never use ether. Keep the applicator in a case or another clean housing.

