

Pevaryl®

For the topical treatment of all forms of dermatomycosis, as well as of microbial mixed infections.

Properties

PEVARYL is a novel antimycotic preparation for topical use. Its range of application includes dermatophytes, yeasts and moulds and therefore covers all significant human pathogenic fungi. Moreover, hitherto therapy resistant mycoses have in numerous cases proved to be curable by PEVARYL. In the presence of mixed infections the antibacterial effect of the preparation offers an additional outstanding advantage.

Active substance

Econazole nitrate 1% = 1-[2-(2,4-dichlorophenyl)-2-(4-chlorbenzylloxy)-ethyl]-imidazol nitrate.

Indications

I. Dermatomycoses caused by

- Dermatophytes

Trichophyton rubrum	
Trichophyton mentogrophytes	
Epidermophyton floccosum	etc.

- Yeasts

Candida albicans	Rhodotorula
Candida guilliermondii	Malassezia furfur
Torulopsis	etc.

Moulds

II. Skin infections caused by gram -positive

Pathogens

- Streptococci
- Staphylococci
- Nocardia minutissima

III. Otitis externa mycotica - mycosis of the auditory canal

Contraindications

None of record

Tolerance and side effects

PEVARYL is readily tolerated even by sensitive skins. Local irritations are extremely infrequent and may occur for example, on eczematous skin areas. In such cases it is advisable to apply an antieczematous therapy a few days prior to treatment with PEVARYL.

Application and dosage

(unless otherwise prescribed by the physician)

PEVARYL is applied to the affected skin regions in the morning and evening and, using a finger, lightly massaged into the skin. Intertriginous areas (e. g., interdigital spaces, gluteal folds) in an oozing state should initially be kept dry with strips of gauze or gauze bandage. In the treatment of onychomycoses the use of occlusive dressings is recommended.

PEVARYL (Spray-) Powder is used to support the PEVARYL Skin Cream, (Spray-) Solution and Skin Lotion therapy, as well as for follow-up and prophylactic treatment. In cases of intertrigo, for example, the application of PEVARYL in powder form alone may be sufficient.

Mycosis of the auditory canal:

Intact ear-drum : Apply 1 - 2 drops of PEVARYL lotion or insert a gauze strip soaked with PEVARYL lotion into the auditory canal.

Regular use of PEVARYL according to instructions is a prerequisite for an enduring cure.

Please store PEVARYL like any other drug beyond the reach of children.

Note

Standard tubes are used by reason of rationalization. Depending on the volume of the charge, a certain vacuum cannot be avoided: however, the net weight corresponds to the regulations in any case.

Commercial forms

- 15 g Skin cream 1%
- 20 g Powder 1%
- 30 ml solution atomizer
- 15 g Lipogel 1%

Dispensed by pharmacies on doctor's prescription.

 Cilag

Manufactured by
MINAPHARM EGYPT

Under Licence of
CILAG - Switzerland

