

Pevisone[®] Cream

For the treatment of inflamed and infected dermatoses.

Composition :

Econazolum nitricum 1% Triamcinolon-acetonidum 0.1% Conserv. (Acid. benzoicum, Butylhydroxyanisolum) Excip. ad unguent emulgat.

Properties :

PEVISONE is a well-balanced combination of econazole nitrate and triamcinolone acetonide, intended for use in dermatology.

Econazole nitrate is a broad-spectrum antimycotic that acts against dermatophytes, yeasts, and moulds; moreover, it is also active against gram positive bacteria (eg. staphylococci).

Triamcinolone acetonide develops marked antiinflammatory, antipruritic, and anti-allergic effects.

Both econazole nitrate and triamcinolone acetonide have been shown to develop their full efficacy in the combination preparation PEVISONE.

PEVISONE cream is based upon an oil-in-water emulsion

PEVISONE ointment is based upon a water-in-oil emulsion

These vehicles make close contact between active substances And skin possible.

Indications :

Eczema: inflammatory dermatoses with mycotic and/or bacterial infection Inflammatory dermatomycoses caused by dermalophytes, yeasts, and moulds.

Mycotic affections in the body folds, also in the presence of bacterial secondary and/or mixed infections.

PEVISONE cream is useful in particular for the treatment of moist . oozing, acute/subacute dermatoses.

PEVISONE ointment is most helpful in the treatment of dry subchronic/chronic dermatoses.

PEVISONE cream and ointment are also indicated for the treatment of:

- Paronychia
- Perleche
- Diaper dermatitis
- Superficial pyoderma.

Tolerability and adverse reactions :

PEVISONE cream and ointment are well tolerated. These preparations are also suitable for therapeutic use on delicate skin. Hypersensitivity was observed in very rare cases only.

Contra-indications :

Like any other preparation containing corticoids PEVISONE is contra-indicated in cases of tuberculous and viral infections (herpes. vaccinia, varicellae) of the skin. It is nowadays generally recommended that in pregnancy, especially during the first trimester, drugs should only be administered after their potential benefits have been thoroughly weighed against possible hazards.

Dosage and application :

(unless prescribed otherwise by the physician)

Apply PEVISONE cream or ointment to the affected skin areas twice a day (morning and evening) and rub it in gently. Like with any other preparation containing corticoids, treatment with PEVISONE should - if possible - be limited to a period of 3 to 4 weeks' in order to prevent skin damage caused by corticoids (atrophy, telangiectases, striae).

Warning: Avoid prolonged application to thin skin and to the face.

Two-week to four-week therapy of eczematous diseases will normally result in complete cure.

Treatment with PEVISONE of inflammatory mycoses should continue until inflammation disappears. If necessary the therapy can then be carried on using PEVARYL until complete cure is obtained.

Warnings :

Like any other drug. PEVISONE cream and ointment should be kept out of the reach of children.

How supplied :

5 g. 15 g. 100 g cream

15 g ointment

At pharmacies, on prescription only.

Manufactured by
MINAPHARM EGYPT
Under Licence of
CILAG - Switzerland

