

Primary Prevention of Acute Respiratory Tract Infections in Children Using a Bacterial Immunostimulant: A Double-Masked, Placebo-Controlled Clinical Trial

Jaime V. Jara-Pérez, MD,¹ and Arturo Berber, MD, PhD²

¹Girl's Home, National Program for the Family, and ²Arzneimittelforschung BASF Pharma, Mexico City, Mexico

ABSTRACT

Background: Acute respiratory tract infections (ARTIs) are among the main causes of morbidity and mortality in children. The bacterial extract OM-85 BV has shown some protective effect for ARTIs in preschool children and a reduction in exacerbations of chronic bronchitis in adults.

Objectives: This trial reports results of a double-masked, placebo-controlled, parallel-group clinical study that assessed the efficacy and tolerability of OM-85 BV in the prevention of ARTIs in school girls living in an orphanage.

Methods: Two hundred girls (age range, 6 to 13 years) living in an orphanage entered the trial. Participants were randomly allocated to receive either OM-85 BV or placebo for 10 consecutive days a month for 3 consecutive months. Patients were followed up for 6 months, including the administration period. The trial began in September 1996 and finished in March 1997. Primary end points were the type and number of infections. Secondary end points included when an infection occurred, time to clinical cure, severity of infection, absenteeism from school due to an ARTI, number of antibiotics or other drugs prescribed, and duration of concomitant drug treatment.

Results: During the trial, patients in the OM-85 BV group experienced 143 ARTIs (135 upper ARTIs and 8 otitis episodes) and patients in the placebo group experienced 299 ARTIs (273 upper ARTIs, 1 lower ARTI, and 25 otitis episodes). The median number of ARTIs was 1.0 (0.0, 3.0; 5th percentile, 95th percentile) in the OM-85 BV group compared with 3.0 (2.0, 4.0; 5th percentile, 95th percentile) in the placebo group. This difference was statistically significant ($P < 0.001$). Participants who received OM-85 BV also showed significantly better results ($P < 0.001$) than participants who received placebo in terms of median duration of illness, median number of missed school days due to an ARTI, median number of antibiotic and drug courses, and median duration of concomitant treatment. There were significant differences ($P < 0.05$) in severity of ARTIs during

Accepted for publication May 3, 2000.

Printed in the USA. Reproduction in whole or part is not permitted.

month 4 of the trial, with patients receiving OM-85 BV showing less severe ARTIs than patients receiving placebo and shorter mean time to clinical cure from the second month to the fourth month. No adverse events related to the trial medications were reported.

Conclusions: OM-85 BV had a preventive effect on ARTIs in the school girls, with a reduction in the antibiotic requirements and the duration of ARTIs. Future studies are needed to further explore the role of OM-85 BV in the prevention of ARTIs.

Key words: acute respiratory tract infection, prevention, immunostimulant, OM-85 BV. (*Clin Ther.* 2000;22:748-759)

INTRODUCTION

By the end of the 20th century, acute respiratory tract infections (ARTIs) were one of the primary causes of childhood morbidity and mortality worldwide.¹ It is estimated that ARTIs have caused 28% of all deaths in children younger than 5 years.¹ ARTIs are responsible for many missed school days² and much parental absenteeism from work.³ They are also often associated with otitis, which is an important cause of hearing loss and poor school performance.⁴

Risk factors for ARTI include attendance at a day care center,⁵ overcrowding at school,^{3,6} having older brothers or sisters,⁶ smoking at home,⁷ and lack of breast-feeding.⁸

Various methods of preventing ARTIs have been studied in clinical trials; for example, general hygiene methods in children attending day care centers⁹; administration of nutritional supplements such as vitamin A to malnourished children,¹⁰ vitamin C in normal and malnourished chil-

dren,¹¹ and trace elements in malnourished and susceptible children¹²; antibiotics¹³; administration of gamma globulins¹⁴; and the use of immunostimulants from various sources, including synthetic,¹⁵ thymic extracts or factors,¹⁶ or biologic, such as *Klebsiella* extracts containing lipopolysaccharide¹⁷ and mixtures of bacterial extracts.¹⁸⁻²¹ Although the therapeutic use of immunostimulants (defined as medications that produce an enhancement of non-specific immunity and an increased infection resistance) is common in some European countries, the efficacy of these medications in the prevention of ARTIs in children is controversial.^{22,23}

OM-85 BV* is an immunostimulant containing the lyophilized extract of the following bacteria: *Haemophilus influenzae*, *Streptococcus pneumoniae*, *Klebsiella pneumoniae*, *Klebsiella ozaenae*, *Staphylococcus aureus*, *Streptococcus pyogenes*, *Streptococcus viridans*, and *Moraxella catarhalis*.²⁴ Previous controlled clinical trials have shown that OM-85 BV reduces the incidence of ARTIs for 6 months in children prone to these infections,²⁵ and for 3 months in children attending day care centers.²⁶ The relative risk for ≥ 4 ARTIs in 6 months was 0.79 (95% confidence interval [CI], 0.59 to 1.06), and a relative risk for ≥ 3 ARTIs in the first 3 months was 0.52 (95% CI, 0.31 to 0.86).²⁶ The protective effects of OM-85 BV in viral ARTI were shown in a trial of children attending day care centers, a study that included viral isolation.²⁷ The protective effects of OM-85 BV have also been shown for exacerbations of chronic bronchitis in adults.^{28,29}

*Trademark: Broncho-Vaxom® (OM PHARMA Meyrin, Geneva, Switzerland; marketed in Mexico by Química Knoll de Mexico).

By the end of 1999, 9387 patients had received OM-85 BV as part of clinical trials. Only 448 (4.8%) had reported adverse events. These were mainly gastrointestinal disturbances (1.7%), exacerbation of symptoms (0.5%), skin disturbances (0.4%), and headache and dizziness (0.3%) (data on file; OM PHARMA Meyrin, Geneva, Switzerland; 1999).

Each capsule of OM-85 BV contains 3.5 mg of the lyophilized extract. The mechanism of action of OM-85 BV is partially known. OM-85 BV is free of lipopolysaccharide, and its effect is mediated by a CD14-independent pathway.³⁰ In macrophage and monocytic cells, OM-85 BV increases intracellular calcium levels and induces the production of glucose-regulated protein³¹ and C-Fos/serum responsive element protein.^{32,33} These second messengers induce the expression of proinflammatory cytokines interleukin (IL) 1- α , IL-6, IL-8, and tissue necrosis factor- α .³²⁻³⁵ In addition, OM-85 BV induces phagocytic cells to produce nitric oxide and oxygen,³⁵ and to express adhesion molecules.^{30,36}

The production of these cytokines and substances may be related to the changes observed in the immunoglobulin (Ig) levels and phagocytic activity in patients receiving OM-85 BV. These patients have shown enhancement of cellular immune responses,^{37,38} and increases of secretory IgA,³⁹⁻⁴¹ serum IgA,^{28,42} and serum IgG and IgM,^{39,42} as well as activation of phagocytic cells.^{40,41}

Because the tolerability and efficacy of OM-85 BV have been studied in children younger than 6 years and adults,²⁵⁻²⁹ we conducted a clinical trial to investigate the tolerability and efficacy of the immunostimulant OM-85 BV in school-age girls.

PATIENTS AND METHODS

This double-masked, placebo-controlled, parallel-group clinical trial began in September 1996 and finished in March 1997. The trial was conducted during the boreal winter because this season had the highest incidence of ARTI, according to the orphanage records. The primary aim of the study was to investigate the effect of OM-85 BV on the incidence of ARTIs. Secondary aims were to investigate the effect of OM-85 BV on absenteeism from school and the use of antibiotics. The girls were informed about the aims and procedures of the trial, and told they could refuse to participate or withdraw their consent at any time, without providing a reason and without being punished or losing any rights.

All participants provided oral consent to participate in the study. In addition, written informed consent was obtained from the legal guardian for each girl (a close relative and if missing, the chairman of the institution).

The protocol and case report forms were approved by the Research and Ethics Committee of the Central Medical Committee of the National Program for the Family (formed by the chairmen and medical directors of the 3 other orphanages in the program) and were in accordance with the Mexican regulations and the Helsinki Declaration of 1975, as revised in 1983.

Based on previous trials in Mexico,²⁰ we expected to have a 50% reduction in the incidence of ARTIs. Using a difference of 1.5 ± 1.5 ARTIs between the groups during 6 months,²⁰ the calculated sample size was 23 patients per group; for a difference of 0.75 ± 1.5 (half of the previously reported efficacy), a sample size of 86 patients per group was needed. An

initial sample size of 100 patients per group was used.

Patients were girls aged 6 to 13 years living in a single orphanage (Girl's Home, National Program for the Family) south of Mexico City. They had been in the orphanage for ≥ 2 years and stayed in 60 bedrooms with 6 beds each, without any special clustering.

Selection criteria were as follows: ≥ 3 ARTIs during the previous 6 months (according to the medical records in the orphanage), negative familial history of allergy, no seasonal or food-related wheezing or nose itching, and no nasal fold. Participants could not have anatomic alterations of the respiratory tract, chronic respiratory diseases (tuberculosis, cystic fibrosis), autoimmune diseases, liver or kidney failure, malnutrition, or cancer, and could not have received treatment with corticosteroids, immunosuppressants, immunostimulants, gamma globulins, or anticonvulsive drugs in the past 6 months. All the girls were between the 60th and 95th percentiles of height and body weight for their respective ages.

There were 450 girls within the age range in the orphanage at the beginning of the trial. We included 200 girls who met the selection criteria and had the highest number of ARTIs.

The names of the participants were sorted alphabetically, and patients were assigned to receive either OM-85 BV or placebo according to a random number list in blocks of 10. The treatment for each patient number was prepared in advance. The boxes, blisters, and capsules for the active treatment and placebo had the same appearance, and the tastes of the powders were similar. A.B. prepared the randomization list and the rest of the materials but did not see the patient list. The cap-

sules were administered by nurses, and the empty blisters were kept to determine compliance (ie, to count the possible missing capsules).

For each patient number there was an opaque sealed envelope with a sheet of paper stating whether the trial medication was OM-85 BV or placebo. All the envelopes were kept available for the investigators in the study center. The envelopes could be opened in the case of serious adverse events. Closed envelopes were recovered by A.B. at the end of the trial.

Patients received 1 capsule orally—OM-85 BV (3.5 mg) or placebo—per day for the first 10 consecutive days of each month for 3 consecutive months. All patients started the trial medication at the same time of day. The trial participants were examined daily by the nurse and medical teams, and the characteristics of ARTIs were recorded on the case report form as they occurred. The following variables were assessed: type and number of infections (main end point), when a girl had an infection, time to clinical cure, severity of the infection, absenteeism (days out of school) due to an ARTI, number of antibiotics or other drugs prescribed, and duration of treatment in days (secondary end points). Severity of infection was assessed using a 114-mm visual analog scale with 3 ratings (very mild, moderate, very severe). Only J.J.-P. marked the scale. In addition, J.J.-P. examined and questioned participants with ARTIs daily up to the end of the episode and completed the case report forms.

An upper ARTI was defined as the presence of ≥ 1 of the following signs: rhinorrhea, sore throat, or cough without signs of a lower ARTI for ≥ 48 hours. A lower ARTI was defined as the presence of ≥ 1 of

the following signs: rales or crepitations, wheezing, stridor, respiratory rate >50 per minute, cyanosis, or chest indrawing (depression of intercostal spaces) for ≥ 48 hours. Otitis was defined as earache with erythema and limited mobility of the tympanic membrane determined by pneumatic otoscopy. Similar upper and lower ARTI definitions have been used in epidemiologic studies in developing countries.⁶ Two infections were counted as such only when the patient was without symptoms for ≥ 72 hours between the end of 1 ARTI and the beginning of another. A treatment course was defined as completion of ≥ 1 day of drug treatment. Clinical cure was defined as complete resolution of all symptoms.

Adverse events were recorded on case report forms as they occurred and reported monthly on an adverse event report form provided with the case report form.

Statistical Analysis

The end point differences between the groups were analyzed using the Mann-Whitney *U* test. In addition, the rate of patients suffering <3 ARTIs throughout the 6-month period were analyzed by Kaplan-Meier statistics defining the event

as the occurrence of 3 ARTIs, and the relative risk for ≥ 1 ARTI, ≥ 2 ARTIs, ≥ 3 ARTIs, and ≥ 1 otitis episode were calculated. The SPSS statistics program (SPSS Inc., Chicago, Illinois) was used for all analyses.

RESULTS

No significant differences were observed between the groups at the beginning of the trial. Baseline demographics are shown in Table I. All participants completed the trial. However, data for 1 girl in the OM-85 BV group were lost. Thus, statistical analysis of results is based on 99 patients who received OM-85 BV and 100 patients who received placebo.

During the trial, patients in the OM-85 BV group had 143 ARTIs (135 upper ARTIs and 8 otitis episodes) and patients in the placebo group had 299 ARTIs (273 upper ARTIs, 1 lower ARTI, and 25 otitis episodes). When the number of ARTIs in each group was compared with the number recorded in clinical files during the 6 months (March 1996 to August 1996) before the trial (494 and 509, respectively), significant differences were detected for both groups ($P < 0.001$ by Mann-Whitney *U* test).

Table I. Baseline patient characteristics.

	OM-85 BV (n = 99)	Placebo (n = 100)
Age (y)*	9.8 \pm 1.9	9.6 \pm 1.9
Body weight (kg)*	32.8 \pm 8.5	33.3 \pm 9.5
Height (cm)*	134.7 \pm 11.6	134.0 \pm 11.9
No. of ARTIs in past 6 months [†]	5 (4;6)	5 (4;6)

ARTIs = acute respiratory tract infections.

*Mean \pm SD.

[†]Median values (percentiles 5;95).




Figure 1. Number of acute respiratory tract infections (ARTIs) by entire group by month. * $P < 0.05$ versus placebo, Mann-Whitney U test.

In our study, the median number of ARTIs was 1.0 (0.0, 3.0; 5th percentile, 95th percentile) in the OM-85 BV group compared with 3.0 (2.0, 4.0; 5th percentile, 95th percentile) in the placebo group. Figure 1 depicts the number of ARTIs by entire group by month. The incidence of ARTI increased in December in both groups, but the number of cases was significantly lower ($P < 0.05$) in the OM-85 BV group.

Table II shows the number and type of ARTIs in each group as well as the median and 5th and 95th percentiles of the trial end points (ie, number of ARTIs, illness duration, number of antibiotic courses, number of drug courses, duration of concomitant treatments, and days out of school). For all variables measured, values were significantly better in the OM-85 BV group than in the placebo group ($P < 0.001$, Mann-Whitney U test). The median dura-

tion of illness was 5.0 days (0.0, 19.0; 5th percentile, 95th percentile) in the OM-85 BV group compared with 19.0 days (12.0, 31.0; 5th percentile, 95th percentile) in the placebo group ($P < 0.001$). The median number of missed school days due to an ARTI was 0.0 days (0.0, 4.6; 5th percentile, 95th percentile) in the OM-85 BV group compared with 3.0 days (0.0, 7.0; 5th percentile, 95th percentile) in the placebo group ($P < 0.001$).

The median numbers of antibiotic and drug courses were 0.0 (0.0, 3.0; 5th percentile, 95th percentile) and 1.0 (0.0, 3.3; 5th percentile, 95th percentile), respectively, in the OM-85 BV group compared with 2.0 (1.0, 4.0; 5th percentile, 95th percentile) and 3.0 (2.0, 4.0; 5th percentile, 95th percentile), respectively, in the placebo group ($P < 0.001$). The median duration of concomitant treatment was 5.0

Table II. Number and type of acute respiratory tract infections (ARTIs) and associated variables during the trial.

	OM-85 BV (n = 99)	Placebo (n = 100)
Total no. of ARTIs	143	299
Upper ARTIs	135	273
Lower ARTIs	0	1
Otitis episodes	8	25
No. of ARTIs*	1.0 (0.0;3.0) [†]	3.0 (2.0;4.0)
Duration of illness (d)*	5.0 (0.0;19.0) [†]	19.0 (12.0;31.0)
No. of antibiotic courses*	0.0 (0.0;3.0) [†]	2.0 (1.0;4.0)
No. of drug courses* [‡]	1.0 (0.0;3.3) [†]	3.0 (2.0;4.0)
Duration of concomitant treatment (d)*	5.0 (0.0;27.1) [†]	24.0 (11.1;35.0)
No. of days out of school* [§]	0.0 (0.0;4.6) [†]	3.0 (0.0;7.0)

*Median values (percentiles 5;95).

[†] $P < 0.001$ versus the placebo group (Mann-Whitney U test for all items).

[‡]Any medication, including antibiotics.

[§]Due to ARTI.

days (0.0, 27.1; 5th percentile, 95th percentile) compared with 24.0 days (11.1, 35.0; 5th percentile, 95th percentile) in the placebo group ($P < 0.001$).

Patients in the OM-85 BV group had shorter time to clinical cure from October through December than the control group (5.66 ± 2.15 vs 6.61 ± 1.56 , 4.89 ± 1.23 vs 6.51 ± 1.56 , and 5.21 ± 1.96 vs 7.89 ± 2.38 , respectively; mean \pm SD; $P < 0.05$, Student t test). No significant difference was observed during the first month, and the numbers of ARTIs in January and February were too small for comparison.

The severity of ARTIs (measured on the 114-mm visual analog scale) was similar in the first 3 months of the trial, but in the fourth month the severity was significantly lower in the OM-85 BV group than in the placebo group (45.13 ± 17.82 vs 64.92 ± 10.66 mm; mean \pm SD; $P < 0.001$, Student t test). The difference be-

tween the 2 groups was -19.79 (95% CI, -25.026 to -14.573). Data on the OM-85 BV group were insufficient for statistical analysis in the fifth and sixth months.

Figure 2 shows the percentage of patients experiencing ≤ 3 ARTIs throughout the 6-month period. The OM-85 BV group had significantly fewer ARTIs than the placebo group ($P < 0.001$, log rank, Breslow and Tarone-Ware tests).

Table III shows the relative risk for suffering ≥ 1 , ≥ 2 , and ≥ 3 ARTIs, and ≥ 1 otitis episode in patients who received OM-85 BV compared with patients who received placebo.

No adverse events related to the use of the trial medications were reported.

DISCUSSION AND CONCLUSIONS

The rationale for the use of immunostimulants for preventing ARTIs is based on




Figure 2. Percentage of patients experiencing ≤ 3 acute respiratory tract infections throughout the 6-month period. * $P < 0.001$ by the log-rank, Breslow and Tarone-Ware tests for the global evolution of the groups; $P < 0.05$ by chi-square test).

the assumption that highly susceptible patients have at least some immune disturbances. It has been shown that patients receiving OM-85 BV have an enhanced cellular immune response, including an increase in secretory IgA; serum IgG, IgM, and IgA; and activation of phagocytic cells.³⁷⁻⁴²

Because most childhood ARTIs have a viral origin,⁴³ it is not clear how a bacterial immunostimulant might prevent these viral infections. The protective effect may be due to the increase in secretory IgA levels, since low secretory IgA levels are associated with higher susceptibility to viral ARTIs.^{44,45} Indeed, the preventive effect of OM-85 BV on viral ARTIs was demonstrated in a multicenter, double-

masked, placebo-controlled clinical trial including viral isolation.²⁷

The retrospective incidence of ARTIs in studied population was 19.29 episodes per 100 child-weeks, a high incidence even in comparison with groups of children younger than 6 years in developing countries (12.7 to 16.8 ARTIs per 100 child-weeks).⁶ In our study, we noticed a high retrospective incidence of ARTIs compared with data reported prospectively in other open-label^{19,20} and double-masked²¹ clinical trials of OM-85 BV. During our trial, the incidence of ARTIs was higher than that found in age-matched, healthy children.⁴⁶ This high incidence could be a reflection of sample selection, overcrowding in the orphanage, long and durable

transmission chains for ARTIs, and exposure to low temperatures during the daily morning shower due to lack of hot water. The reduction in incidence in the placebo group and to a lesser degree in the OM-85 BV group could be ascribed to a previous overreport, an improvement in the general measures for ARTI prevention (wearing coats and avoiding exposure to low temperatures during the morning shower [generally believed to be the main predisposing factors]), possibly to the partial interruption of ARTI transmission by OM-85 BV administration, differences in seasonal rates of ARTI, or simply a result of chance. Because the trial was conducted in boreal fall and winter, the incidence reduction cannot be related to seasonal temperature change.

We tried to exclude patients with the clinically suspected allergies (familial history of allergy, seasonal or food-related wheezing or nose itching, or nasal fold), yet the presence of allergies cannot be completely ruled out. As the definitions for upper and lower ARTIs overlap with allergy symptoms, it is possible that in some cases, allergies were diagnosed as ARTIs. However, because the 2 groups were well balanced, the possible effect of allergies could be regarded as similar in both groups.

We considered the study population to have no immunologic defects because they did not have any chronic or serious conditions and were of normal height and body weight. Thus, we regarded the study population as a group of healthy girls overexposed to ARTI pathogens.

In our trial, OM-85 BV reduced the incidence of ARTIs significantly (a median reduction of 2 episodes), and this reduction resulted in shorter duration of ARTIs (median reduction of 14 days), less concomitant drug consumption (median re-

Table III. Relative risk for acute respiratory tract infections (ARTIs) and otitis episodes in the OM-85 BV group compared with the placebo group.

	Relative Risk	95% Confidence Interval
≥1 ARTI	0.889	0.829 – 0.953
≥2 ARTIs	0.381	0.295 – 0.493
≥3 ARTIs	0.194	0.117 – 0.319
≥1 otitis	0.333	0.157 – 0.706

duction of 19 days of treatment), and lower incidence of otitis (a reduction of 66%). All these issues may have an important pharmaco-economic impact, particularly in day care centers and elementary schools. It was not possible to assess the effect of OM-85 BV on lower ARTIs, because almost all the infections that occurred in the 2 groups were upper ARTIs.

The favorable therapeutic results and good tolerance are in accordance with results obtained in preschool children,^{25,26} as well as in secondary prevention studies in preschool and school children aged 6 to 12 years.¹⁸⁻²¹

No adverse events were associated with the study medication. Such a low adverse-event incidence has been reported in previous Mexican trials.¹⁸⁻²¹ The global incidence of adverse events in OM-85 BV clinical trials is 4.8%, comprising mainly gastrointestinal and skin disturbances, exacerbation of the symptoms, and headache and dizziness (data on file, OM PHARMA Meyrin, Geneva, Switzerland, 1999).

This nonspecific approach to the prevention of ARTIs should be further explored as a new alternative to provide protection against ARTIs.

ACKNOWLEDGMENTS

The trial medications and case report forms were provided by Química Knoll de México SA de CV BASF Pharma, Mexico City, Mexico.

The authors thank Ms. Marit Meredith and Ms. Lucila Velasco for reviewing the manuscript.

Address correspondence to: Arturo Berber, MD, Arzneimittelforschung BASF Pharma, La Candelaria 186, Colonia Atlantida, Mexico City D.F. C.P. 04370, Mexico.

REFERENCES

1. Campbell H. Acute respiratory infection: A global challenge. *Arch Dis Child*. 1995; 73:281–286.
2. Haskins R, Kotch J. Day care and illness: Evidence, costs, and public policy. *Pediatrics*. 1986;77(Suppl 2):951–982.
3. Bell DM, Gleiber DW, Mercer AA, et al. Illness associated with child day care: A study of incidence and cost. *Am J Public Health*. 1989;79:479–484.
4. Erwick-Roberts J, Burchinal MR, Collier AM, et al. Otitis media in early childhood and cognitive, academic and classroom performance of the school-aged child. *Pediatrics*. 1989;83:477–485.
5. Schwartz B, Giebink GS, Henderson FW, et al. Respiratory infections in day care. *Pediatrics*. 1994;94(6 Pt 2):1018–1020.
6. Selwyn BJ. The epidemiology of acute respiratory tract infection in young children: Comparison of findings from several developing countries. *Rev Infect Dis*. 1990;12(Suppl 8):S870–S888.
7. Jin C, MacKay-Rossignol A. Effects of passive smoking on respiratory illness from birth to age eighteen months, in Shanghai, People's Republic of China. *J Pediatr*. 1993;123:553–558.
8. Wright AL, Holberg CJ, Martinez FD, et al. Breast feeding and lower respiratory tract illness in the first year of life. *Br Med J*. 1989;299:946–949.
9. The Child Day Care Infectious Disease Study Group. Public health considerations of infectious diseases in child day care centers. *J Pediatr*. 1984;105:683–701.
10. Barreto ML, Santos LM, Assis AM, et al. Effect of vitamin A supplementation on diarrhoea and acute lower-respiratory-tract infections in young children in Brazil. *Lancet*. 1994;344:228–231.
11. Hemila H. Vitamin C intake and susceptibility to the common cold. *Br J Nutr*. 1997;77:59–72.
12. Sazawal S, Black RE, Jalla S, et al. Zinc supplementation reduces the incidence of acute lower respiratory infections in infants and preschool children: A double-blind, controlled trial. *Pediatrics*. 1998; 102:1–5.
13. Dajani A, Taubert K, Ferrieri P, et al. Treatment of acute streptococcal pharyngitis and prevention of rheumatic fever. A statement for health professionals. Committee on Rheumatic Fever, Endocarditis, and Kawasaki Disease of the Council on Cardiovascular Disease in the Young, the American Heart Association. *Pediatrics*. 1995;96(4 Pt 1):758–764.
14. Nydahl-Persson K, Petterson A, Fasth A. A prospective, double-blind, placebo-controlled trial of i.v. immunoglobulin and trimethoprim-sulfamethoxazole in children with recurrent respiratory tract infections. *Acta Paediatr*. 1995;84:1007–1009.

15. Passali D, Calearo C, Conticello S. Pidotimod in the management of recurrent pharyngotonsillar infections in childhood. *Arzneim-Forsch Drug Res.* 1994;44:1511-1516.
16. De Mattia D, Decandia P, Ferrante P, et al. Effectiveness of thymostimulin and study of lymphocyte-dependent antibacterial activity in children with recurrent respiratory infections. *Immunopharmacol Immunotoxicol.* 1993;15:447-459.
17. Dahan R, Constantini D, Caulin C, Yagello M. Clinical trials with C 1740, an immunomodulator compound proposed for prevention of acute infectious exacerbations in chronic bronchitis. *Methods Find Exp Clin Pharmacol.* 1986;8:41-43.
18. Berber AC, Del-Rio-Navarro BE. Use of Broncho-Vaxom in private practice: Phase IV trial in 587 children. *Clin Ther.* 1996; 18:1068-1079.
19. Field J, Gómez-Barreto D, Del-Rio-Navarro BE, Berber A. Use of OM-85 BV in the primary prevention of acute respiratory tract infections in children in orphanages. *Curr Ther Res Clin Exp.* 1998;59: 407-418.
20. Gutiérrez-Tarango MD, Berber A. Efficacy of a bacterial extract (OM-85 BV) in preventing recurrent respiratory tract infections in susceptible children. *Clin Drug Invest.* 1997;13:76-84.
21. Gómez-Barreto D, De la Torre C, Álvarez A, et al. Safety and efficacy of OM-85-BV plus amoxicillin/clavulanate in the treatment of subacute sinusitis and the prevention of recurrent infections in children. *Allergol Immunopathol.* 1998;26:17-22.
22. Valleron AJ, Grimfeld A. Evaluation of clinical trials of immunomodulators for prevention of recurrent respiratory infections in children. *Dev Biol Stand.* 1992; 77:149-158.
23. Collet JP. Immunomodulators and primary prevention of respiratory infections: Methodological considerations. *Dev Biol Stand.* 1992;77:159-165.
24. Bessler WG, Sedelmeir E. Biological activity of bacterial cell wall components. *Arzneim-Forsch Drug Res.* 1993;43:502-507.
25. Paupe J. Immunotherapy with an oral bacterial extract (OM-85 BV) for upper respiratory infections. *Respiration.* 1991;58: 150-154.
26. Collet JP, Ducruet T, Kramer MS, et al. Stimulation of nonspecific immunity to reduce the risk of recurrent infections in children attending day-care centers. *Pediatr Infect Dis J.* 1993;12:648-652.
27. Aymard M, Chomel JJ, Allard JP, et al. Epidemiology of viral infections and evaluation of the potential benefit of OM-85 BV on the virologic status of children attending day-care centers. *Respiration.* 1994; 61(Suppl 1):24-31.
28. Cvoriscec B, Ustar M, Pardon R, et al. Oral immunotherapy of chronic bronchitis: A double-blind placebo-controlled multicentre study. *Respiration.* 1989;55:129-135.
29. Collet JP, Shapiro S, Ernst P, et al. Effects of an immunostimulating agent on acute exacerbations and hospitalizations in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* 1997; 156:1719-1724.
30. Marchant A, Goldman M. OM-85 BV up-regulates the expression of adhesion molecules on phagocytes through a CD14-independent pathway. *Int J Immunopharmacol.* 1996;18:259-262.
31. Baladi S, Kantengwa S, Donati YRA, Polla BS. Effects of the bacterial extract OM-85 on phagocyte functions and the stress response. *Med Inflamm.* 1994;3:143-148.

32. Keul R, Roth M, Papakonstantinou E, et al. Induction of interleukin 6 and interleukin 8 expression by Broncho-Vaxom (OM-85 BV) via C-Fos/serum responsive element. *Thorax*. 1996;51:150-154.
33. Roth M, Keul R, Papakonstantinou E, et al. Characterization of intracellular signaling transduction and transcription factors involved in Broncho-Vaxom (OM-85 BV)-induced expression of interleukin-6 and interleukin-8 in human pulmonary fibroblasts. *Eur Respir Rev*. 1996;6:171-175.
34. Broug-Holub E, Persoons JHA, Schornagel K, Kraal G. Changes in cytokine and nitric oxide secretion by rat alveolar macrophages after oral administration of bacterial extracts. *Clin Exp Immunol*. 1995;101:302-307.
35. Broug-Holub E, Kraal G. Cytokine production by alveolar macrophages after oral administration of OM-85 BV in a rat model. *Eur Respir Rev*. 1996;6:163-165.
36. Jacquier-Sarlin MR, Dreher D, Polla BS. Selective induction of the glucose-regulated protein grp78 in human monocytes by bacterial extract (OM-85): A role for calcium as second messenger. *Biochem Biophys Res Commun*. 1996;226:166-171.
37. Mavël J. Stimulation of immunoprotective mechanisms by OM-85 BV: A review of results from in vivo and in vitro studies. *Respiration*. 1994;61(Suppl 1):8-15.
38. Maestroni GJ, Losa GA. Clinical and immunobiological effects of an orally administered bacterial extract. *Int J Immunopharmacol*. 1984;6:111-117.
39. Puigdollers JM, Serna GR, Hernández del Rey I, et al. Immunoglobulin production in man stimulated by orally administered bacterial lysate. *Respiration*. 1980;40:142-149.
40. Emmerich B, Emslander HP, Pachmann K, et al. Local immunity in patients with chronic bronchitis and the effects of a bacterial extract, Broncho-Vaxom, on T lymphocytes, macrophages, gamma-interferon and secretory immunoglobulin A in bronchoalveolar lavage fluid and other variables. *Respiration*. 1990;57:90-99.
41. Lusuardi M, Capelli A, Carli S, et al. Local airways immune modifications induced by oral bacterial extracts in chronic bronchitis. *Chest*. 1993;103:1783-1791.
42. Djuric O, Mihailovic-Vucinic V, Stojcic V. Effect of Broncho-Vaxom on clinical and immunological parameters in patients with chronic obstructive bronchitis. A double-blind, placebo-controlled study. *Int J Immunother*. 1989;5:139-143.
43. Gadomski AM. Potential interventions for preventing pneumonia among young children: Lack of effect of antibiotic treatment for upper respiratory infections. *Pediatr Infect Dis J*. 1993;12:115-120.
44. Jemmott JB, McClelland DC. Secretory IgA as a measure of resistance to infectious disease: Comments on Stone, Cox, Valdimarsdottir and Neale. *Behav Med*. 1989;15:63-71.
45. Rossen RD, Butler WT, Waldman RH, et al. The proteins in nasal secretion II. A longitudinal study of IgA and neutralizing antibody levels in nasal washing from men infected with influenza virus. *JAMA*. 1970;211:1157-1161.
46. Monto AS, Ullman BM. Acute respiratory illness in an American community: The Tecumseh study. *JAMA*. 1974;227:164-169.