

Lucidril 250 mg

COMPANY NAME : Minapharm- Egypt

TRADE NAME : Lucidril 250 F.C tablets.

GENERIC NAME : Meclofenoxate Hydrochloride 250 mg

COMPOSITION

Compostion Per F.C. Tablets:

Active Ingredient:

Meclofenoxate Hydrochloride (freeze-dried).....250 mg

Inactive Ingredient:

Tribasic Calcium Phosphate.....95 mg
Magnesium Stearate.....5 mg
Polyvinyl acetate.....20 mg
Methyl Cellulose.....3.5 mg
Ethyl Cellulose.....0.5 mg
Dibutyl phthalate.....0.75 mg
Talc.....25 mg

PHARMACEUTICAL FORM:

Film coated tablets.

PHARMACOLOGICAL ACTIONS:

-Has antianoxic effect whether anoxia is provoked by defect or misuse of oxygen.

-Improves extraction and utilization of glucose from the brain which allows maintenance of storage of energy for the neuron even in case of O₂ deficiency or aging. This action is also mediated by stimulation of aerobic glycolysis or via the enhancement of anaerobic respiration.

This metabolic activity is accompanied with cerebral vasodilatation which is more marked in the grey matter in human being.

-Oppose the deposition of aging pigments in the cerebral cells.

-Have different effects on the cerebral cells as it passes the blood brain barrier.

INDICATIONS:

- Cerebral Aging, Cerebral atherosclerosis, neurological or psychological sequelae of head injury; manifesting by: mental deterioration; mental fatigue & inability to concentrate, disturbed memory and mood, intellectual impairment and behavioral changes. Neurological symptoms; Transient neurological attacks, paresis of face and limbs, Visual disturbance, Vertigo & tinnitus, Motor & sensory aphasia.

DOSAGE:

2 tablets in the morning, 2 tablets at noon (before 4 O'clock p.m.) in repeated courses, not less than one month.

The degree of action of Lucidril is very largely dependent upon the duration of treatment.

CONTRA-INDICATION:

History of allergy to any of the ingredients of the drug.
Pronounced condition of excitement.

SIDE EFFECTS :

Rarely allergic reactions.

DRUG-DRUG INTERACTION :

Must not be administered along with medications containing Citicoline.

PREGNANCY & LACTATION :

As a general rule if you are pregnant or breast feeding you should always seek the advice of your doctor or pharmacist before taking a medication.

PRECAUTION & WARNING :

Athletes can be careful: this product contains an active ingredient capable of inducing a positive dope test result.

PACKAGE:

Carton box of 20 tablets, each containing 250 mg of meclufenoxate hydrochloride.

STORAGE:

Lucidril should be stored at temperature not exceeding 30°C.

**Manufactured by: Minapharm Co. for
Pharmaceuticals & Chemical Industries**