

110 mm

# *Lucidril 500*

**Meclofenoxate**

## **COMPOSITION:**

Each tablet contains Meclofenoxate Hydrochloride 500mg.

## **INDICATIONS :**

1. Manifestations of cerebral insufficiency:
  - Disturbance of memory and attention
  - Vertigo
  - Ringing and humming in the ears
  - Visual disturbance
  - Personality disorders
  - Intellectual deterioration
2. Sequelae of cerebrovascular accidents, hemiplegia and aphasia.
3. Head injury and its sequelae: headache, vertigo, personality disorders, memory disturbance and aphasic disorders.

## **DOSE AND ADMINISTRATION:**

To be taken as directed by the doctor.

Usual doses: 2 tablets per day (last dose before 4 PM).

## **PRECAUTIONS AND WARNINGS:**

Athletes caution: this product contains an active ingredient which can induce a positive reaction to the tests made as a part of antidoping controls.

Owing to a slight stimulant effect, it is advisable to avoid administration of this drug at the end of the day.

## **CONTRA-INDICATIONS:**

History of allergy to any of the ingredients of the drug.  
Pronounced conditions of excitement.

## **DRUG-DRUG INTERACTIONS:**

No interaction is known up to now.

## **PREGNANCY AND LACTATION:**

As a general rule if you are pregnant or breast feeding you should always seek the advice of your doctor or pharmacist before taking a medication.

## **STORAGE:**

Store below 30°C.

## **PRESENTATION:**

It is supplied in 2 strips of 10 tablets.

Manufactured by  
**MINAPHARM** Egypt



180 mm