

SRILANE 5 %, cream

Idroclamide

IDENTIFICATION OF THE MEDICINAL PRODUCT

Composition: Each tube contains 20 g cream

Idroclamide (INN)	per 100 g
Excipient: incl. propylene glycol	5 g
Emulgade F® , octyldodecanol, purified water q.s. for	10 g
	100 g

Pharmacotherapeutic class

- OTHER TOPICAL DRUG FOR JOINT AND MUSCLE PAINS (M: Locomotor system)

MERCK SANTE - 37, rue Saint Romain - 69008 LYON

Manufacturer: MERCK SANTE - 115, av Lacassagne - 69003 LYON

IN WHICH CASE(S) CAN THIS DRUG BE USED?

This drug is recommended, in adults and children over 15 years of age, as local adjunct short-term treatment in:

- benign joint injuries of the common ankle sprain type and contusions.
- acute lower back pain.

WARNING!

When should this drug not be used?

This drug MUST NOT BE USED in the following cases:

- history of allergy to one of the components of the cream,
- injured skin, regardless of the lesion: weeping lesions, eczema, infected lesions, burns or open wounds,
- children under 15 years of age.

If you are in any doubt, you must consult your doctor or your pharmacist for advice.

Special warnings

Do not apply to the mucous membranes or the eyes.

Precautions for use

The hands must be washed after use to avoid any involuntary contact of the product with the eyes or mucous membranes.

If in doubt do not hesitate to seek the advice of your doctor or pharmacist.

Drug interactions and other interactions

In order to avoid possible interactions between various drugs, you must always tell your doctor or your pharmacist about any other treatment you are receiving.

Pregnancy- breastfeeding

As a general rule, if you are pregnant or breastfeeding you should always seek the advice of your doctor or pharmacist before taking a medication.

List of excipients whose knowledge is necessary for risk-free use in some patients

- each 100 g of cream contains 10 g of propylene glycol.

HOW SHOULD THIS DRUG BE USED?

Dosage, method and route of administration

Cutaneous route - RESTRICTED TO ADULTS AND CHILDREN OVER 15 YEARS OF AGE. Apply as local massages to the painful area, 1 to 3 times per day, until completely rubbed in. Wash hands after use.

Frequency and time at which the drug should be taken

Do not apply more than 3 times per day.

Duration of treatment

Do not use for a prolonged period without medical advice.

UNDESIRABLE AND UNPLEASANT EFFECTS

Like all active substances, this drug may, in some individuals, cause more or less unpleasant effects:

- possible local allergic reaction requiring discontinuation of treatment,
- possibility of normal redness of the skin at the site of application.

Report any undesirable or unpleasant effect not mentioned in this leaflet to your pharmacist or doctor.

STORAGE

Do not use past the expiry date indicated in clear on the package.

DATE OF LAST LEAFLET REVISION: December 1998

MINAPHARM

Under Licence of
Merck Santé France

MERCK
SERONO



MINAPHARM

سريلان 5% كريم

إيدروسيلاميد

بيان التركيب: كل أنبوبة تحتوي على 20 جم كريم لكل 100 جم

إيدروسيلاميد 5 جم

مواد مألوفة بها جليكول 10 جم

إملجيد، أوكتيل دوديكانول وماء كمية كافية حتى 100 جم

دواعى الإستعمال:

يوصف هذا الدواء للبالغين وللأطفال فوق سن 15 عاماً كمعالج موضعى قصير المدى فى الحالات التالية:

- آلام إصابة إلتواء المفاصل أو الكدمات.

- فى حالات آلام الظهر السفلى.

تحذيرات:

- لا يستعمل هذا الدواء فى حالات الحساسية لمواد الكريم.

- لا يستعمل فى حالات الجروح والحروق والإكزيما والالتهابات الجلدية.

- لا يستعمل للأطفال تحت سن 15 عاماً.

- يحذر من إستعماله فى العين ولذلك يجب غسل اليدين جيداً بعد إستعماله المقتن.

- لتفادى أى تفاعل لهذا الدواء مع دواء آخر إستشر طبيبك قبل الإستخدام، وكذلك فى حالات الحمل

والرضاعة.

طريقة الإستعمال:

ضع الكريم على المكان المصاب وذلك جيداً حتى يتم إمتصاصه من خلال الجلد ولا يستعمل أكثر من ثلاث

مرات فى اليوم.

الأثار الجانبية:

قد يحدث حساسية توجب وقف الإستعمال وقد يحدث إحمرار فى الجلد فى منطقة الدهان.

إنتاج **مينا فارم** مصر

بنصريح من شركة
ميرك سانتيه فرنسا



MINAPHARM

SRILANE 5%
11 X 18 cm.

P. No.: 286